



From traditional Gurmar to modern phytomedicine: Traditional knowledge, phytochemistry and emerging applications of *Gymnema sylvestre*

Dhale D A

Department of Botany, SSVPS's, L.K.Dr.P.R. Ghogrey Science College, Dhule, Maharashtra, India

Abstract

Gymnema sylvestre (Retz.) R.Br. ex Sm., commonly known as Gurmar or Madhunashini, is an important medicinal climber with a long history of use in Indian traditional medicine. Its characteristic ability to temporarily suppress sweet taste has earned it the popular name 'sugar destroyer.' Traditionally, its leaves and other parts have been used for glycosuria, diabetes-related disorders, digestive and urinary ailments, inflammation, and other conditions. Modern research has linked these traditional uses with diverse bioactive constituents, particularly gymnemic acids, gymnemasaponins, gymnemosides, and gurmardin. Gymnemic acids interact with the human sweet-taste receptor complex TAS1R2/TAS1R3, providing a molecular basis for sweetness suppression. Experimental studies have further reported effects on intestinal glucose absorption, insulin secretion, glucose utilization, oxidative stress, lipid metabolism, and inflammatory pathways. Early clinical studies of standardized leaf extracts reported improvements in glycemic parameters, while subsequent randomized controlled research has provided additional evidence supporting further clinical investigation. However, clinical evidence remains limited by small sample sizes, variation in extract composition, dosage, treatment duration, and inadequate standardization. Consequently, modern applications of *G. sylvestre* extend beyond its traditional antidiabetic use to include functional foods, natural taste-modifying agents, standardized botanical preparations, nutraceuticals, metabolic-health research, and biotechnological production of bioactive compounds. This review examines the transition of *G. sylvestre* from traditional medicine to modern applications, focusing on its ethnobotany, phytochemistry, molecular mechanisms, pharmacological and clinical evidence, safety, biotechnology, conservation, and future research prospects.

Keywords: *Gymnema sylvestre*, gurmar, madhunashini, gymnemic acids, gurmardin, diabetes, sweet taste suppression

Introduction

Medicinal plants represent an important interface between traditional knowledge and contemporary drug discovery. Numerous modern pharmacological investigations have originated from observations accumulated through traditional healthcare systems. *Gymnema sylvestre* is a particularly interesting example because its traditional reputation as a "sugar destroyer" is supported by a distinctive sensory phenomenon as well as by experimental and clinical investigations.

G. sylvestre is currently accepted botanically as a member of the family Apocynaceae and is native across tropical and subtropical regions of the Old World, including India and several regions of tropical Asia and Africa (Royal Botanic Gardens, Kew, 2026) [18]. It is a perennial woody climber and is particularly associated with the Indian medicinal-plant tradition. The plant is known by several vernacular names, including Gurmar, Madhunashini, Meshashringi, Kavali and related regional names.

The medicinal importance of *G. sylvestre* is primarily associated with its leaves, which have a long history of use in traditional systems of medicine, particularly for the management of glycosuria, diabetes, metabolic and urinary disorders. The plant, popularly known as Gurmar or Madhunashini, has also been traditionally employed for various digestive, respiratory, inflammatory and other health-related conditions, reflecting its broad ethnomedicinal significance (Tiwari *et al.*, 2014; Di Fabio *et al.*, 2015) [4, 23]. The traditional use of *G. sylvestre* for disorders associated with abnormal glucose metabolism has subsequently stimulated extensive experimental and clinical

research. Pharmacological investigations have demonstrated that its leaf constituents may influence several aspects of glucose homeostasis, including intestinal glucose absorption, insulin secretion, glucose utilization and pancreatic β -cell function (Baskaran *et al.*, 1990; Imoto, 1990; Persaud *et al.*, 1999; Tiwari *et al.*, 2014; Khan *et al.*, 2019; Devangan *et al.*, 2021) [2, 3, 8, 10, 15, 23].

The transition from traditional medicine to modern phytomedicine has been facilitated by the identification of gymnemic acids and related triterpenoid constituents. These compounds provide a biochemical explanation for the temporary suppression of sweetness and have stimulated research into their possible applications in metabolic health, food science and taste modification. Molecular studies have demonstrated that gymnemic acids interact with the human sweet receptor TAS1R2/TAS1R3, particularly involving the transmembrane domain of TAS1R3 (Sanematsu *et al.*, 2014) [19].

Thus, *G. sylvestre* represents a useful model for understanding how ethnobotanical knowledge can progress through phytochemical characterization, mechanistic pharmacology, clinical research and modern product development.

Botanical Identity and Distribution

1. Taxonomic position

Kingdom: Plantae

Order: Gentianales

Family: Apocynaceae

Genus: *Gymnema*

Species: *Gymnema sylvestre* (Retz.) R.Br. ex Sm.

The species was historically placed within Asclepiadaceae, a group now generally treated within Apocynaceae. The accepted name according to the current Kew taxonomic database is *Gymnema sylvestre* (Retz.) R.Br. ex Sm. It is a perennial, woody, scandent climber or twiner, characterized by white latex and a well-developed taproot. The young shoots are shortly villous, while older stems become woody and lenticellate. Leaves are simple, opposite and shortly petiolate, with blades broadly ovate, obovate, elliptic or suborbicular; the apex is acute to shortly attenuate, the base rounded or weakly cordate, and both surfaces may be pubescent. The plant commonly grows in tropical and subtropical forests, scrublands and thickets (Tiwari *et al.*, 2014; Khan *et al.*, 2019; Royal Botanic Gardens, Kew, 2026) ^[10, 18, 23].



Fig 1: Habitat of *Gymnema sylvestre*

The inflorescences are extra-axillary, usually forming dense, subumbelliform clusters that are shorter than the adjacent leaves. Flowers are small, bisexual and pentamerous, generally greenish-white to pale yellow, with a pubescent calyx and a campanulate, five-lobed corolla bearing a characteristic corona. The fruit is typically a paired, elongated follicle containing numerous seeds, each provided with a silky coma that facilitates wind dispersal. The opposite leaves, milky latex, compact flower clusters and paired follicles together provide useful diagnostic characters for identification of *G. sylvestre*, while the leaves remain the principal medicinally important organ (Kanetkar *et al.*, 2007; Tiwari *et al.*, 2014; Khan *et al.*, 2019) ^[9, 10, 23].

2. Distribution

The species has a broad tropical and subtropical distribution extending through the Indian subcontinent, Southeast Asia, parts of Africa and Australia. In India, it is associated particularly with peninsular and central regions and occurs in forest and scrub habitats.

The plant is a woody climber with opposite leaves, small flowers and elongated follicles. The leaves are the principal medicinally utilized part and constitute the major commercial source of gymnemic-acid-containing preparations (Kirtikar & Basu, 1987) ^[11].

Traditional Knowledge and Ethnomedicinal Significance

The traditional significance of *G. sylvestre* is particularly strong in Ayurveda and Indian ethnomedicine. The plant is popularly associated with the Sanskrit-derived concept of Madhunashini, meaning “destroyer of sweetness,” reflecting its characteristic effect on sweet taste.

Historical Ayurvedic literature and subsequent ethnobotanical documentation associate the plant with glycosuria, urinary disorders and metabolic conditions. Traditional applications have also included dyspepsia, constipation, jaundice, hemorrhoids, respiratory disorders and other ailments. Different communities in India have used various parts of the plant according to local medicinal traditions (Tiwari *et al.*, 2014; Di Fabio *et al.*, 2015) ^[4, 23].

The ethnobotanical importance of *G. sylvestre* is significant for two reasons. First, traditional use identified a reproducible physiological phenomenon-suppression of sweetness-that could subsequently be investigated experimentally. Second, its long association with glycemic disorders provided a rationale for pharmacological research into glucose metabolism.

The transition can therefore be represented as:

Traditional observation → ethnobotanical documentation → phytochemical isolation → molecular mechanism → pharmacological validation → clinical investigation → standardized phytoproduct development

This pathway provides a valuable model for the rational development of medicinal plants.

Phytochemical Profile

The major bioactive constituents of *G. sylvestre* include gymnemic acids, gymnemasaponins, gymnemosides and the peptide gurmarin. Gymnemic acids are structurally diverse oleanane-type triterpenoid saponins, and individual congeners differ considerably in their antisweet activity (Yoshikawa *et al.*, 1991, 1992; Murakami *et al.*, 1996; Tiwari *et al.*, 2014; Khan *et al.*, 2019) ^[10, 14, 23, 25, 26]. More recent sensory-guided investigations have demonstrated substantial differences among individual gymnemic acids, with some compounds exhibiting particularly strong inhibition of human sweet-receptor responses (Pickrahn *et al.*, 2021; Manisha *et al.*, 2025) ^[7, 13].

1. Gymnemic acids

Gymnemic acids are the best-known constituents of *G. sylvestre*. They are triterpenoid glycosides and occur as a mixture of related compounds rather than as one chemically homogeneous molecule. Structural diversity among gymnemic acids is associated with differences in glycosylation and acylation patterns.

Gymnemic acids are principally responsible for the characteristic antisweet activity of the plant. Their ability to inhibit sweet taste has made them important not only in pharmacology but also in sensory science and food research (Sanematsu *et al.*, 2014) ^[16].

2. Gymnemasaponins and gymnemosides

The leaves contain additional triterpenoid saponins, including gymnemasaponins and gymnemosides. These compounds contribute to the broader biological activity reported for *G. sylvestre* extracts.

3. Gurmarin

Gurmarin is a peptide associated with the antisweet activity of *G. sylvestre*. Together with gymnemic acids, it has contributed to the development of the plant as a model for studying taste perception.

4. Other constituents

Phytochemical investigations have reported flavonoids, anthraquinone-related compounds, sterols, resins and other secondary metabolites in different parts of the plant. Consequently, whole extracts may exhibit biological activities that cannot necessarily be attributed to gymnemic acids alone (Tiwari *et al.*, 2014) ^[23].

From “Sugar Destroyer” to Molecular Mechanism

One of the most remarkable properties of *G. sylvestre* is its ability to temporarily reduce the perception of sweetness after exposure of the oral cavity to its leaf-derived constituents.

Modern sensory biology has established that sweet taste is primarily mediated through the heterodimeric G-protein-coupled receptor TAS1R2/TAS1R3. Experimental studies demonstrated that gymnemic acids suppress responses mediated by the human sweet receptor and identified an important role for the transmembrane domain of TAS1R3 (Sanematsu *et al.*, 2014) ^[19].

This discovery provides a striking example of how a traditional sensory observation can be translated into molecular pharmacology.

The mechanism can be summarized as:

Gymnemic acids → interaction with TAS1R2/TAS1R3 → reduced sweet-receptor signaling → temporary reduction in perceived sweetness

Importantly, this does not mean that *G. sylvestre* eliminates the physiological need for glucose or that it can independently treat diabetes. Rather, the finding demonstrates a molecular basis for one specific sensory property of the plant.

A systematic review of human studies on plant-derived sweet-taste suppressors concluded that evidence regarding downstream effects on food consumption remains inconsistent and that additional research is needed to determine whether acute sweetness suppression produces meaningful long-term changes in eating behavior (Rayo Morales *et al.*, 2023) ^[17].

Antidiabetic and Metabolic Potential

The antidiabetic potential of *G. sylvestre* has received considerably more attention than any other pharmacological property.

1. Modulation of glucose absorption

Experimental research suggests that gymnemic acids and related constituents can interfere with intestinal glucose absorption. Earlier physiological investigations demonstrated inhibitory effects on glucose absorption, providing one possible explanation for reduced postprandial glycemic responses (Imoto, 1990; Grover *et al.*, 2002) ^[6, 8].

2. Effects on insulin secretion

Experimental evidence also indicates that *G. sylvestre* extracts can influence pancreatic β -cell function. Persaud *et*

al. (1999) ^[15] reported stimulation of insulin release *in vitro* from pancreatic preparations and β -cell models.

These findings suggest that the plant may influence glucose homeostasis through multiple mechanisms rather than through a single target.

3. Potential effects on glucose utilization

Experimental literature has proposed effects on glucose-utilizing enzymes, hepatic glucose metabolism and other metabolic pathways. These mechanisms remain more firmly established in experimental systems than in large-scale human trials and should therefore be interpreted cautiously (Tiwari *et al.*, 2014) ^[23].

4. Lipid metabolism

Animal and preliminary human research has also suggested possible hypolipidemic effects. Changes in cholesterol and triglyceride metabolism have been reported in experimental studies and in some early clinical investigations, although stronger controlled clinical evidence is needed before definitive therapeutic claims can be made (Kumari, 2024) ^[12].

Clinical Evidence

The clinical history of *G. sylvestre* illustrates both the promise and limitations of herbal medicine research.

One of the earliest frequently cited clinical investigations was conducted by Baskaran *et al.* (1990) ^[2], in which a water-soluble leaf extract designated GS4 was administered to patients with non-insulin-dependent diabetes mellitus in addition to conventional treatment. The investigators reported reductions in blood glucose and glycosylated indices and observed that some participants required lower doses of conventional antidiabetic medicines.

Another early investigation evaluated GS4 in patients receiving insulin therapy and reported reductions in fasting glucose, HbA1c and insulin requirements (Shanmugasundaram *et al.*, 1990) ^[2].

These studies were historically important but had methodological limitations. A systematic review of herbal treatments for glycemic control subsequently characterized the evidence for *G. sylvestre* as suggestive but inconclusive, particularly because of the limited number and quality of clinical trials (Yeh *et al.*, 2003) ^[24].

More recently, Gaytán Martínez *et al.* (2021) ^[5] conducted a randomized, double-blind, placebo-controlled clinical trial involving 30 individuals with impaired glucose tolerance. Participants received *G. sylvestre* or placebo, with metabolic and insulin-related outcomes assessed before and after intervention. The study represents an important step toward modern clinical evaluation, although the relatively small sample size means that larger and longer trials remain necessary.

Table 1: Representative evidence supporting the traditional-to-modern transition of *Gymnema sylvestre*

Research stage	Major observation	Significance
Traditional medicine	Use for glycosuria and metabolic disorders	Ethnopharmacological foundation
Traditional sensory observation	Reduction of sweetness after chewing leaves	Identified a distinctive biological effect
Phytochemistry	Isolation and characterization of gymnemic acids	Linked activity to chemical constituents
Molecular biology	Interaction with TAS1R2/TAS1R3	Mechanistic explanation of antisweet activity

Experimental pharmacology	Effects on glucose absorption and insulin secretion	Multiple possible metabolic mechanisms
Early clinical research	Improvements in glycemic parameters reported with GS4	Initial human evidence
Modern clinical research	Randomized placebo-controlled study in impaired glucose tolerance	Progress toward evidence-based validation
Product development	Extracts, teas, tablets, supplements and functional formulations	Translation into contemporary applications

Antioxidant and Anti-inflammatory Potential

Oxidative stress and chronic low-grade inflammation are important components of several metabolic disorders. *G. sylvestre* extracts contain phenolic and other antioxidant-associated constituents and have demonstrated free-radical-scavenging and anti-inflammatory effects in experimental studies.

However, the presence of antioxidant activity in chemical or cellular assays should not automatically be interpreted as evidence that consumption of the plant prevents oxidative-stress-related disease in humans. Further studies should identify the active constituents, pharmacokinetic behavior and clinically relevant concentrations.

The broader pharmacological literature has reported antioxidant, anti-inflammatory, antimicrobial, hepatoprotective and other activities, but the strength of evidence varies substantially among endpoints (Di Fabio *et al.*, 2015; Tiwari *et al.*, 2014) [4, 23].

Antiobesity and Appetite-Related Applications

The antisweet property of *G. sylvestre* has generated considerable interest in weight-management research. The rationale is that temporary suppression of sweet taste may modify the sensory reward associated with highly sweet foods.

This concept has potential applications in the development of taste-modifying foods and dietary products. Nevertheless, the relationship between short-term taste suppression and sustained reduction in energy intake remains uncertain. A 2023^[17] systematic review concluded that human evidence regarding the downstream effects of sweet-taste suppression on eating behavior is inconsistent and requires further investigation (Rayo Morales *et al.*, 2023) [17].

Therefore, *G. sylvestre* should not be presented as an established weight-loss treatment. Its more scientifically defensible modern application is as a source of taste-modifying bioactive compounds with potential relevance to dietary behavior and food formulation.

Functional Foods and Nutraceutical Applications

The distinctive sensory properties of *G. sylvestre* have opened opportunities beyond conventional herbal medicine. Commercial and experimental applications include:

- Herbal teas and infusions.
- Standardized leaf extracts.
- Tablets and capsules.
- Functional beverages.
- Nutraceutical formulations.
- Taste-modifying preparations.
- Research ingredients for reduced-sugar food design.
- Botanical combinations targeting metabolic health.

Tiwari *et al.* (2014) [23] noted that *G. sylvestre* preparations have been incorporated into tea bags, health tablets, supplements, beverages and confectionery-related products. The modern food-industry opportunity lies particularly in the possibility of using gymnemic-acid-containing preparations as sensory modulators, rather than treating the

plant exclusively as an antidiabetic herb (Pothuraju, *et al.*, 2014) [16].

Modern Pharmacological Applications Beyond Diabetes

Although glycemic regulation remains the primary research focus, experimental studies have investigated several other activities.

1. Antimicrobial activity

Extracts and isolated constituents have demonstrated antimicrobial activity against selected microorganisms. This supports the traditional use of medicinal plants for infectious and inflammatory conditions, but clinical translation requires pathogen-specific and formulation-specific evidence.

2. Hepatoprotective potential

Experimental investigations have suggested protective effects against selected forms of chemically induced hepatic injury. Such observations warrant further mechanistic research but should not be interpreted as proof of treatment efficacy in human liver disease.

3. Cardiovascular and lipid-related potential

Hypolipidemic and antiatherosclerotic effects have been reported in animal models. For example, experimental work has investigated the effect of leaf extracts on lipid profiles and atherosclerotic processes. These findings provide a rationale for additional research into metabolic syndrome and cardiovascular risk but do not establish clinical efficacy.

4. Anticancer research

Several phytochemicals from *G. sylvestre* have been investigated experimentally for antiproliferative and cytotoxic effects. Such work is valuable for drug-discovery research, but these findings remain predominantly preclinical and should not be equated with an established anticancer treatment.

Biotechnology and Sustainable Production

The increasing demand for *Gymnema sylvestre* as a medicinal and nutraceutical plant highlights the need for sustainable production systems that can ensure a continuous supply of quality raw material while reducing pressure on natural populations. Conventional propagation may be insufficient to meet commercial requirements, particularly when wild populations are affected by habitat degradation, overharvesting and environmental variability. In this context, plant biotechnology offers important opportunities for both conservation and large-scale production. *In vitro* micropropagation and tissue-culture techniques have demonstrated the potential for rapid multiplication of *G. sylvestre* and the production of relatively uniform planting material, thereby supporting the establishment of cultivated populations and conservation programmes (Ahmed *et al.*, 2009; Tiwari *et al.*, 2014; Khan *et al.*, 2019) [1, 10, 23]. Such approaches can also facilitate the selection and multiplication of elite plants with desirable growth characteristics and higher concentrations of pharmacologically important metabolites.

Advanced biotechnological approaches, including callus culture, organ culture, cell-suspension culture, hairy-root culture and elicitation, may further provide opportunities for enhancing the production of gymnemic acids and other valuable secondary metabolites under controlled conditions. Metabolite profiling combined with molecular authentication and DNA barcoding can improve the identification of genuine plant material, while controlled cultivation and standardized harvesting can minimize variations associated with genotype, geographical origin, soil conditions, climate, plant developmental stage and post-harvest processing (Kanetkar *et al.*, 2007; Tiwari *et al.*, 2014; Khan *et al.*, 2019) [9, 10, 23]. Therefore, an integrated strategy involving *in vitro* propagation, conservation of genetic resources, elite-chemotype selection, controlled cultivation and phytochemical standardization could provide a sustainable pathway for meeting the growing demand for *G. sylvestre* while supporting the long-term conservation of natural populations. Recent Indian initiatives in medicinal-plant conservation have similarly emphasized the importance of seed propagation, vegetative multiplication and *in vitro* techniques for conserving medicinal plant diversity threatened by habitat loss and unsustainable harvesting.

Standardization and Quality Control

Standardization and quality control are essential for translating traditional *G. sylvestre* preparations into reliable modern phytopharmaceuticals and nutraceutical products. The expression “*G. sylvestre* extract” may represent preparations with substantially different phytochemical compositions because the concentration and profile of bioactive constituents are influenced by geographical origin, genetic variability, plant part, stage and season of harvesting, extraction solvent and temperature, processing conditions, storage, and the concentration and composition of gymnemic acids and other triterpenoid saponins (Kanetkar *et al.*, 2007; Tiwari *et al.*, 2014; Khan *et al.*, 2019) [9, 10, 23]. Therefore, botanical authentication of the raw material, documentation of a voucher specimen, clearly defined extraction procedures, extract yield, and quantitative estimation of characteristic marker compounds are important prerequisites for reproducible research and product development. Standardization should additionally include chromatographic fingerprinting, assessment of microbial contamination, heavy metals and pesticide residues, together with appropriate stability studies to ensure the identity, purity, potency and safety of the final preparation (Tiwari *et al.*, 2014; Di Fabio *et al.*, 2015; Khan *et al.*, 2019) [4, 10, 23].

Modern analytical techniques can significantly improve the quality assessment of *G. sylvestre* preparations. High-performance liquid chromatography (HPLC) and liquid chromatography–mass spectrometry (LC–MS) can be employed for qualitative and quantitative characterization of gymnemic acids and related saponins, while chromatographic fingerprinting provides a useful approach for detecting batch-to-batch variations and possible adulteration. Advanced metabolomic approaches can further characterize the complex chemical profile of the plant and help establish relationships between specific phytochemical constituents and biological activities (Tiwari *et al.*, 2014; Khan *et al.*, 2019) [10, 23]. Consequently, the development of validated analytical methods, authenticated plant material,

standardized extraction protocols and chemically defined marker profiles is crucial for ensuring reproducibility, efficacy and safety and for facilitating the transition of *G. sylvestre* from traditional herbal medicine to evidence-based phytopharmaceutical and nutraceutical applications.

Future Research Perspectives

The future of *G. sylvestre* research lies in moving from broad claims of “antidiabetic activity” toward precise, mechanism-based and clinically validated applications.

Important research priorities include:

1. Standardized clinical trials

Large, multicenter, randomized controlled trials should evaluate standardized extracts with clearly defined gymnemic-acid concentrations.

2. Precision phytochemistry

Advanced metabolomics should be used to identify chemical markers responsible for specific biological effects.

3. Molecular target identification

Further investigation of TAS1R2/TAS1R3 and intestinal glucose-transport pathways may enable development of novel taste-modifying and metabolic-health agents.

4. Gut–metabolic interactions

The interaction between *G. sylvestre* constituents, intestinal glucose handling, gut microbiota and metabolic signaling represents an emerging research area.

5. Functional food design

Gymnemic acids may have potential as natural taste-modifying ingredients in reduced-sugar formulations. However, sensory effectiveness, stability, palatability and regulatory acceptability require systematic study.

6. Biotechnology

Cell culture, elicitation and metabolic engineering could potentially provide controlled production of valuable gymnemic acids and related metabolites.

7. Conservation genomics

Molecular characterization of natural populations could support identification of elite chemotypes and conservation of genetic diversity.

Traditional Knowledge as a Platform for Modern Drug Discovery

The history of *Gymnema sylvestre* provides a compelling example of how traditional medicinal knowledge can serve as a starting point for modern drug discovery and evidence-based phytopharmacology. Rather than being regarded either as definitive therapeutic evidence or as merely anecdotal information, traditional observations can be considered hypothesis-generating biological evidence that requires systematic botanical, phytochemical, pharmacological and clinical validation. In traditional Indian medicine, *G. sylvestre* has long been associated with the management of glycosuria and metabolic disorders, while the characteristic ability of its leaves to temporarily suppress the perception of sweetness provided an important sensory observation that subsequently attracted scientific investigation (Kanetkar *et al.*, 2007; Tiwari *et al.*, 2014; Di

Fabio *et al.*, 2015) [4, 9, 23]. Phytochemical investigations subsequently identified gymnemic acids, gymnemasaponins, gymnemosides and the peptide gurmarin as important bioactive constituents, thereby providing a chemical basis for several of the plant's traditional properties (Yoshikawa *et al.*, 1991, 1992; Murakami *et al.*, 1996; Tiwari *et al.*, 2014) [14, 23, 25, 26].

The progression from traditional knowledge to modern application has subsequently involved several interconnected stages: ethnomedicinal observation → botanical and ethnobotanical validation → phytochemical characterization → molecular investigation → experimental pharmacology → clinical evaluation → standardized product development. Molecular studies have demonstrated that gymnemic acids can interact with the human sweet-taste receptor system, particularly TAS1R2/TAS1R3, providing a mechanistic explanation for the traditional observation of sweetness suppression (Sanematsu *et al.*, 2014) [19]. Experimental investigations have further explored effects on intestinal glucose absorption, insulin secretion, pancreatic β-cell function and glucose metabolism (Imoto, 1990; Persaud *et al.*, 1999; Tiwari *et al.*, 2014; Khan *et al.*, 2019) [8, 10, 15, 23]. Early clinical studies subsequently investigated standardized *G. sylvestre* preparations in individuals with diabetes and reported potentially beneficial effects on glycemic parameters, although the available clinical evidence remains limited and heterogeneous (Baskaran *et al.*, 1990; Shanmugasundaram *et al.*, 1990; Devangan *et al.*, 2021) [2, 3]. This progression illustrates how traditional knowledge can guide modern pharmacological discovery while emphasizing that traditional claims must ultimately be supported by reproducible phytochemical characterization, mechanistic evidence, controlled clinical trials, safety evaluation and quality standardization. Consequently, *G. sylvestre* has evolved from a traditionally recognized “sugar-destroying” medicinal plant into a scientifically investigated source of bioactive compounds with potential applications in standardized botanical medicines, nutraceuticals, functional foods, metabolic-health research and natural taste-modifying technologies.

Conclusion

Gymnema sylvestre represents one of the most compelling examples of a medicinal plant progressing from traditional knowledge toward modern phytomedicine. Its historical reputation as Gurmar or Madhunashini is closely connected with the distinctive ability of its leaf constituents to suppress sweet taste. Modern molecular research has demonstrated that gymnemic acids interact with the human sweet-taste receptor system, providing a mechanistic explanation for an observation recognized in traditional practice centuries ago. Beyond taste modification, experimental and early clinical research indicates potential effects on glucose absorption, insulin secretion, glucose metabolism, lipid metabolism, oxidative stress and inflammation. However, the evidence should be interpreted within the limitations of existing research, particularly small clinical samples, inconsistent extract standardization and the shortage of large, well-controlled long-term clinical trials.

The future significance of *G. sylvestre* may therefore extend beyond its conventional description as an “antidiabetic herb.” Its constituents represent potential leads for natural taste modifiers, metabolic-health products, functional foods, nutraceuticals and standardized botanical preparations. At

the same time, biotechnology, molecular authentication and sustainable cultivation can help ensure reliable raw-material supply and conservation of genetic resources.

The most productive modern approach is not to replace traditional knowledge with laboratory science, but to integrate the two. Traditional observations can identify promising biological hypotheses; phytochemistry can identify candidate molecules; molecular biology can reveal mechanisms; pharmacology can establish biological activity; and rigorous clinical trials can determine therapeutic relevance. *G. sylvestre* thus provides a valuable model for the transition from traditional medicinal knowledge to evidence-based modern applications.

References

1. Ahmed ABA, Rao AS, Rao MV, T M. In vitro production of gymnemic acid from *Gymnema sylvestre* (Retz.) R.Br. ex Roemer and Schultes through callus culture under stress conditions. In Book, Protocols for in vitro cultures and secondary metabolite analysis of medicinal plants, 2009. DOI: 10.1007/978-1-60327-287-2_8
2. Baskaran K, Ahamath BK, Shanmugasundaram KR, Shanmugasundaram ERB. Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin-dependent diabetes mellitus patients. Journal of Ethnopharmacology, 1990;30(3):295–305. [https://doi.org/10.1016/0378-8741\(90\)90108-6](https://doi.org/10.1016/0378-8741(90)90108-6)
3. Devangan S, Varghese B, Johny E, Gurram S, Adela R. The effect of *Gymnema sylvestre* supplementation on glycemic control in type 2 diabetes patients: A systematic review and meta-analysis. Phytotherapy Research, 2021;35(12):6802–6812. <https://doi.org/10.1002/ptr.7265>
4. Di Fabio G, Romanucci V, Di Marino C, Pisanti A, Zarrelli A. *Gymnema sylvestre* R. Br., an Indian medicinal herb: Traditional uses, chemical composition, and biological activity. Current Pharmaceutical Biotechnology, 2015;16(6):506–516. <https://doi.org/10.2174/138920101606150407112903>
5. Gaytán Martínez LA, Sánchez-Ruiz LA, Zúñiga LY, González-Ortiz M, Martínez-Abundis E. Effect of *Gymnema sylvestre* administration on glycemic control, insulin secretion, and insulin sensitivity in patients with impaired glucose tolerance. Journal of Medicinal Food, 2021, 24(1). <https://doi.org/10.1089/jmf.2020.0024>
6. Grover JK, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. Journal of Ethnopharmacology, 2002;81(1):81–100. [https://doi.org/10.1016/S0378-8741\(02\)00059-4](https://doi.org/10.1016/S0378-8741(02)00059-4)
7. Pickrahn S, Dawid C, Babinger T, Schmid C, Brockhoff A, Meyerhof W, et al. Sensory-guided multidimensional exploration of antisweet principles from *Gymnema sylvestre* (Retz.) Schult. Journal of Agricultural and Food Chemistry, 2021;69(19):5510–5527. doi:10.1021/acs.jafc.1c00994
8. Imoto T. Sugar discriminations and plant-originated glycosides, gymnemic acids. Journal of Biophysics, 1990;30(3):146–150. <https://doi.org/10.2142/biophys.30.146>
9. Kanetkar P, Singhal R, Kamat M. *Gymnema sylvestre*: A memoir. Journal of Clinical Biochemistry and

- Nutrition,2007;41(2):77–81. <https://doi.org/10.3164/jcbtn.2007010>
10. Khan F, Sarker MMR, Ming LC, Mohamed IN, Zhao C, Sheikh BY, et al. Comprehensive review on phytochemicals, pharmacological and clinical potentials of *Gymnema sylvestre*. *Frontiers in Pharmacology*,2019;10:1223. <https://doi.org/10.3389/fphar.2019.01223>
 11. Kirtikar KR, Basu BD. Indian medicinal plants (2nd ed.). Bishen Singh Mahendra Pal Singh, 1987.
 12. Kumari V. Ethnopharmacological importance of *Gymnema sylvestre*. *European Journal of Medicinal Plants*,2024;35(6):224–233. <https://doi.org/10.9734/ejmp/2024/v35i61221>
 13. Manisha M, Bihani M, Khan S, Dang R, Bhat D. Phytochemical and pharmacological insights of *Gymnema sylvestre*: A review. *International Journal of Pharmacognosy and Life Sciences*,2025;6(2):235–240. <https://doi.org/10.33545/27072827.2025.v6.i2c.185>
 14. Murakami N, Murakami T, Kadoya M, Matsuda H, Yamahara J, Yoshikawa M. New hypoglycemic constituents in “gymnemic acid” from *Gymnema sylvestre*. *Chemical and Pharmaceutical Bulletin*,1996;44(2):469–471. <https://doi.org/10.1248/cpb.44.469>
 15. Persaud SJ, Al-Majed H, Raman A, Jones PM. *Gymnema sylvestre* stimulates insulin release in vitro by increased membrane permeability. *Journal of Endocrinology*,1999;163(2):207–212. <https://doi.org/10.1677/joe.0.1630207>
 16. Pothuraju R, Sharma RK, Chagalamarri J, Jangra S, Kavadi PK. A systematic review of *Gymnema sylvestre* in obesity and diabetes management. *Journal of the Science of Food and Agriculture*,2014;94(5):834–840. <https://doi.org/10.1002/jsfa.6458>
 17. Rayo Morales R, Garcia-Burgos D, et al. Suppression of sweet taste-related responses by plant-derived bioactive compounds and eating. Part I: A systematic review in humans. *Food Quality and Preference*, 2023.
 18. Royal Botanic Gardens, Kew. *Gymnema sylvestre* (Retz.) R.Br. ex Sm. *Plants of the World Online*, 2026.
 19. Sanematsu K, Kusakabe Y, Shigemura N, Hirokawa T, Nakamura S, Imoto T, et al. Molecular mechanisms for sweet-suppressing effect of gymnemic acids. *Journal of Biological Chemistry*,2014;289(37):25711–25720. <https://doi.org/10.1074/jbc.M114.560409>
 20. Shanmugasundaram ERB, Gopinath KL, Shanmugasundaram KR, Rajendran VM. Possible regeneration of the islets of Langerhans in streptozotocin-diabetic rats given *Gymnema sylvestre* leaf extracts. *Journal of Ethnopharmacology*,1990;30(3):265–279. [https://doi.org/10.1016/0378-8741\(90\)90106-4](https://doi.org/10.1016/0378-8741(90)90106-4)
 21. Shanmugasundaram ERB, Rajeswari G, Baskaran K, Rajesh Kumar BR, Shanmugasundaram KR, Ahmath BK. Use of *Gymnema sylvestre* leaf extract in the control of blood glucose in insulin-dependent diabetes mellitus. *Journal of Ethnopharmacology*,1990;30(3):281–294. [https://doi.org/10.1016/0378-8741\(90\)90107-5](https://doi.org/10.1016/0378-8741(90)90107-5)
 22. Shiyovich A, Sztarkier I, Nesher L. Toxic hepatitis induced by *Gymnema sylvestre*, a natural remedy for type 2 diabetes mellitus. *American Journal of the Medical Sciences*,2010;340(6):514–517. <https://doi.org/10.1097/MAJ.0b013e3181f41168>
 23. Tiwari P, Mishra BN, Sangwan NS. Phytochemical and pharmacological properties of *Gymnema sylvestre*: An important medicinal plant. *BioMed Research International*,2014;2014:Article 830285. <https://doi.org/10.1155/2014/830285>
 24. Yeh GY, Eisenberg DM, Kaptchuk TJ, Phillips RS. Systematic review of herbs and dietary supplements for glycemic control in diabetes. *Diabetes Care*,2003;26(4):1277–1294. <https://doi.org/10.2337/diacare.26.4.1277>
 25. Yoshikawa K, Arihara S, Matsuura K. A new type of antisweet principles occurring in *Gymnema sylvestre*. *Tetrahedron Letters*,1991;32(6):789–792. [https://doi.org/10.1016/S0040-4039\(00\)74887-8](https://doi.org/10.1016/S0040-4039(00)74887-8)
 26. Yoshikawa K, Nakagawa M, Yamamoto R, Arihara S, Matsuura K. Antisweet natural products. V. Structures of gymnemic acids VIII–XII from *Gymnema sylvestre* R.Br. *Chemical and Pharmaceutical Bulletin*,1992;40(7):1779–1782. <https://doi.org/10.1248/cpb.40.1779>